arXiv:math/0302015v1 [math.CO] 2 Feb 2003

A NOTE ON SUM OF k-TH POWER OF HORADAM'S SEQUENCE

Toufik Mansour [1]

Department of Mathematics, Chalmers University of Technology, S-412 96 Göteborg, Sweden

`toufik@math.chalmers.se`

Abstract

Let $w_{n+2} = pw_{n+1} + qw_n$ for $n \geq 0$ with $w_0 = a$ and $w_1 = b$. In this paper we find an explicit expression, in terms of determinants, for $\sum_{n\geq 0} w_n^k x^n$ for any $k \geq 1$. As a consequence, we derive all the previously known results for this kind of problems, as well as many new results.

Key words: Horadam's sequence, Fibonacci numbers, Lucas numbers, Pell numbers

1. Introduction and the Main result

As usual, Fibonacci numbers F_n, Lucas numbers L_n, and Pell numbers P_n are defined by the second-order linear recurrence sequence

$$w_{n+2} = pw_{n+1} + qw_n, \tag{1}$$

with given $w_0 = a, w_1 = b$ and $n \geq 0$. This sequence was introduced, in 1965, by Horadam [Ho], and it generalizes many sequences (see [HW, HM]). Examples of such sequences are Fibonacci numbers sequence $(F_n)_{n\geq 0}$, Lucas numbers sequence $(L_n)_{n\geq 0}$, and Pell numbers sequence $(P_n)_{n\geq 0}$, when one has $p = q = b = 1$, $a = 0$; $p = q = b = 1$, $a = 2$; and $p = 2$, $q = b = 1$, $a = 0$; respectively. In this paper we interested in studding the generating function for powers of Horadam's sequence, that is, $\mathcal{H}_k(x; a, b, p, q) = \mathcal{H}_k(x) = \sum_{n\geq 0} w_n^k x^n$.

In 1962, Riordan [R] found the generating function for powers of Fibonacci number. He proved the generating function $\mathcal{F}_k(x) = \sum_{n\geq 0} F_n^k(x)$ that satisfy the recurrence relation

$$(1 - a_k x + (-1)^k x^2)\mathcal{F}_k(x) = 1 + kx \sum_{j=1}^{[k/2]} (-1)^j \frac{a_{kj}}{j} \mathcal{F}_{k-2j}((-1)^i x)$$

for $k \geq 1$, where $a_1 = 1$, $a_2 = 2$, $a_s = a_{s-1} + a_{s-2}$ for $s \geq 3$, and $(1 - x - x^2)^{-j} = \sum a_{kj} x^{k-2j}$. Recently, Haukkanen [Ha] studied linear combinations of Horadam's sequences and the generating function of the ordinary product of two of Horadam's sequences. The main result of this paper can be formulated as follows.

---

[1] Research financed by EC's IHRP Programme, within the Research Training Network "Algebraic Combinatorics in Europe", grant HPRN-CT-2001-00272

Let $\Delta_k = \Delta_k(p,q)$ be the $k \times k$ matrix

$$\begin{pmatrix}
1-p^kx-q^kx^2 & -xp^{k-1}q^1\tbinom{k}{1} & -xp^{k-2}q^2\tbinom{k}{2} & \cdots & -xp^2q^{k-2}\tbinom{k}{k-2} & -xpq^{k-1}\tbinom{k}{k-1} \\
-p^{k-1}x & 1-xp^{k-2}q^1\tbinom{k-1}{1} & -xp^{k-3}q^2\tbinom{k-1}{2} & \cdots & -xpq^{k-2}\tbinom{k-1}{k-2} & -xq^{k-1}\tbinom{k-1}{k-1} \\
-p^{k-2}x & -xp^{k-3}q^1\tbinom{k-2}{1} & 1-xp^{k-4}q^2\tbinom{k-2}{2} & \cdots & -xq^{k-2}\tbinom{k-2}{k-2} & 0 \\
-p^{k-3}x & -xp^{k-4}q^1\tbinom{k-3}{1} & -xp^{k-5}q^2\tbinom{k-3}{2} & \cdots & 0 & 0 \\
\vdots & \vdots & \vdots & & \vdots & \vdots \\
-p^2x & -xpq^1\tbinom{2}{1} & -xq^2\tbinom{2}{2} & \cdots & 1 & 0 \\
-p^1x & -xq^1\tbinom{1}{1} & 0 & \cdots & 0 & 1
\end{pmatrix},$$

and let $\delta_k = \delta_k(p,q,a,b)$ be the $k \times k$ matrix

$$\begin{pmatrix}
a^k+g_kx & -xp^{k-1}q^1\tbinom{k}{1} & -xp^{k-2}q^2\tbinom{k}{2} & \cdots & -xp^2q^{k-2}\tbinom{k}{k-2} & -xpq^{k-1}\tbinom{k}{k-1} \\
g_{k-1}x & 1-xp^{k-2}q^1\tbinom{k-1}{1} & -xp^{k-3}q^2\tbinom{k-1}{2} & \cdots & -xpq^{k-2}\tbinom{k-1}{k-2} & -xq^{k-1}\tbinom{k-1}{k-1} \\
g_{k-2}x & -xp^{k-3}q^1\tbinom{k-2}{1} & 1-xp^{k-4}q^2\tbinom{k-2}{2} & \cdots & -xq^{k-2}\tbinom{k-2}{k-2} & 0 \\
g_{k-3}x & -xp^{k-4}q^1\tbinom{k-3}{1} & -xp^{k-5}q^2\tbinom{k-3}{2} & \cdots & 0 & 0 \\
\vdots & \vdots & \vdots & & \vdots & \vdots \\
g_2x & -xpq^1\tbinom{2}{1} & -xq^2\tbinom{2}{2} & \cdots & 1 & 0 \\
g_1x & -xq^1\tbinom{1}{1} & 0 & \cdots & 0 & 1
\end{pmatrix},$$

where $g_j = (b^j - a^jp^j)a^{k-j}$ for all $j = 1, 2, \ldots, k$.

Theorem 1.1. *The generating function $\mathcal{H}_k(x)$ is given by* $\dfrac{\det(\delta_k)}{\det(\Delta_k)}$.

The paper is organized as follows. In Section 2 we give the proof of Theorem 1.1 and in Section 3 we give some applications for Theorem 1.1.

2. PROOFS

Let $(w_n)_{n\geq 0}$ be a sequence satisfied Relation (1) and k be any positive integer. We define a family $\{A_{k,d}\}_{d=1}^{k}$ of generating functions by

$$A_{k,d}(x) = \sum_{n\geq 0} w_n^{k-d} w_{n+1}^{d} x^{n+1}. \tag{2}$$

Now we introduce two relations (Lemma 2.1 and Lemma 2.2) between the generating functions $A_{k,d}(x)$ and $\mathcal{H}_k(x)$ that play the crucial roles in the proof of Theorem 1.1.

Lemma 2.1. *For any* $k \geq 1$,

$$(1 - p^k x - q^k x^2)\mathcal{H}_k(x) - x \sum_{j=1}^{k-1} \binom{k}{j} p^{k-j} q^j A_{k,k-j}(x) = a^k + x(b^k - a^k p^k).$$

Proof. Using the binomial theorem (see [WR]) we get

$$w_{n+2}^k = (pw_{n+1} + qw_n)^k = p^k w_{n+1}^k + \sum_{j=1}^{k-1} \binom{k}{j} p^{k-j} q^j w_{n+1}^{k-j} w_n^j + q^k w_n^k.$$

Multiplying by x^{n+2} and summing over all $n \geq 0$ with using Definition (2) we have

$$\mathcal{H}_k(x) - b^k x - a^k = p^k x(\mathcal{H}_k(x) - a^k) + x \sum_{j=1}^{k-1} \binom{k}{j} p^{k-j} q^j A_{k,k-j}(x) + q^k x^2 \mathcal{H}_k(x),$$

as requested. ☐

Lemma 2.2. *For any* $k - 1 \geq d \geq 1$,

$$A_{k,d}(x) - a^{k-d} b^d x = p^d x(\mathcal{H}_k(x) - a^k) + x \sum_{j=1}^{d} \binom{d}{j} p^{d-j} q^j A_{k,k-j}(x).$$

Proof. Using the binomial theorem (see [WR]) we have

$$w_n^{k-d} w_{n+1}^d = w_n^{k-d}(pw_n + qw_{n-1})^d = w_n^{k-d} \sum_{j=0}^{d} \binom{d}{j} p^{d-j} q^j w_n^{d-j} w_{n-1}^j.$$

Multiplying by x^{n+1} and summing over all $n \geq 0$ we get

$$A_{k,d}(x) - a^{k-d} b^d x = p^d x(\mathcal{H}_k(x) - a^k) + x \sum_{j=1}^{d} \binom{d}{j} p^{k-j} q^j A_{k,k-j}(x),$$

as requested ☐

Proof. (Theorem 1.1) By using the above lemmas together with definitions we get

$$\Delta_k \cdot [\mathcal{H}_k(x), A_{k,k-1}(x), A_{k,k-2}(x), \ldots, A_{k,1}(x)]^{\mathrm{T}} = v_k,$$

where

$$v_k = \left[a^k + x(b^k - a^k p^k),\ (a^1 b^{k-1} - p^{k-1} a^k)x,\ (a^2 b^{k-2} - p^{k-2} a^k)x, \ldots,\ (a^{k-1} b^1 - p^1 x a^k)x\right]^{\mathrm{T}}.$$

Hence, the solution of the above equation gives the generating function $\mathcal{H}_k(x) = \frac{\det(\delta_k)}{\det(\Delta_k)}$, as claimed in Theorem 1.1. ☐

3. Applications

In this section we present some applications for Theorem 1.1.

3.1. **Fibonacci numbers.** If $a = 0$ and $p = q = b = 1$, then Theorem 1.1 for $k = 1, 2, 3, 4, 5, 6$ yields Table 1.

k	The generating function $\mathcal{H}_k(x;1,1,1,1)$	$\mathcal{H}_k(1/100)$
1	$\frac{x}{1-x-x^2}$	$\frac{100}{9899}$
2	$\frac{x(1-x)}{(1+x)(1-3x+x^2)}$	$\frac{9900}{979801}$
3	$\frac{x(1-2x-x^2)}{(1+x-x^2)(1-4x-x^2)}$	$\frac{979900}{96940301}$
4	$\frac{x(1+x)(1-5x+x^2)}{(1-x)(1+3x+x^2)(1-7x+x^2)}$	$\frac{31986700}{3161716833}$
5	$\frac{x(1-7x-16x^2+7x^3+x^4)}{(1-x-x^2)(1+4x-x^2)(1-11x-x^2)}$	$\frac{9284070100}{916060399199}$
6	$\frac{x(1-x)(1-11x-64x^2-11x^3+x^4)}{(1+x)(1-3x+x^2)(1+7x+x^2)(1-18x+x^2)}$	$\frac{97194791100}{9554028773189}$

Table 1. The generating function for the powers of Fibonacci numbers

k	The generating function $\mathcal{H}_k(x;2,1,1,1)$	$\mathcal{H}_k(1/100)$
1	$\frac{2-x}{1-x-x^2}$	$\frac{19900}{9899}$
2	$\frac{4-3x-5x^2}{(1+x)(1-3x+x^2)}$	$\frac{3969500}{979801}$
3	$\frac{8-5x-36x^2+7x^3}{(1+x-x^2)(1-4x-x^2)}$	$\frac{794640700}{96940301}$
4	$\frac{16-15x-180x^2+156x^3+17x^4}{(1-x)(1+3x+x^2)(1-7x+x^2)}$	$\frac{52773853900}{3161716833}$
5	$\frac{32-45x-835x^2+1440x^3+745x^4-31x^5}{(1-x-x^2)(1+4x-x^2)(1-11x-x^2)}$	$\frac{31467947446900}{916060399199}$
6	$\frac{64-167x-3708x^2+12323x^3+12597x^4-3188x^5-65x^6}{(1+x)(1-3x+x^2)(1+7x+x^2)(1-18x+x^2)}$	$\frac{688573873901500}{9554028773189}$

Table 2. The generating function for the powers of Lucas numbers

k	The generating function $\mathcal{H}_k(x;0,1,2,1)$
1	$\frac{x}{1-2x-x^2}$
2	$\frac{x(1-x)}{(1+x)(1-6x+x^2)}$
3	$\frac{x(1-4x-x^2)}{(1+2x-x^2)(1-14x-x^2)}$
4	$\frac{x(1+x)(1-14x+x^2)}{(1-x)(1+6x+x^2)(1-34x-x^2)}$
5	$\frac{x(1-38x-130x^2+38x^3+x^4)}{(1-2x-x^2)(1-82x-x^2)(1+14x-x^2)}$
6	$\frac{x(x-1)(1-104x-1210x^2-104x^3+x^4)}{(1+x)(1+34x+x^2)(1-6x+x^2)(1-198x+x^2)}$

Table 3. The generating function for the powers of Pell numbers

3.2. **Lucas numbers.** If $a = 2$ and $p = q = b = 1$, then Theorem 1.1 for $k = 1, 2, 3, 4, 5, 6$ yields Table 2.

3.3. **Pell numbers.** If $a = 0$, $b = q = 1$ and $p = 2$, then Theorem 1.1 for $k = 1, 2, 3, 4, 5, 6$ yields Table 3.

3.4. **Chebyshev polynomials of the second kind.** If $a = 1$, $b = p = 2t$ and $q = -1$, then Theorem 1.1 for $k = 1, 2, 3, 4, 5, 6$ yields Table 4.

More generally, if applying Theorem 1.1 for $k = 1, 2, 3, 4$, then we get the following corollary.

Corollary 3.1. *Let $k = 1, 2, 3, 4$. Then the generating function $\mathcal{H}_k(x)$ is given by $\frac{\mathcal{A}_k(x)}{\mathcal{B}_k(x)}$ where*

k	The generating function $\mathcal{H}_k(x;1,2t,2t,-1)$
1	$\frac{1}{1-2tx+x^2}$
2	$\frac{1+x}{(1-x)((1+x)^2-4xt^2)}$
3	$\frac{1+4tx+x^2}{(1-2tx+x^2)(1+2t(3-4t^2)x+x^2)}$
4	$\frac{(1+x)((1-x)^2+12t^2x)}{(1-x)((1+x)^2-4t^2x)(16t^2(1-t^2)x+(1-x)^2))}$
5	$\frac{1-6tx+2x^2+32t^3x+96t^4x^2+32t^3x^3-32t^2x^2-6x^3t+x^4}{(1+2t(3-4t^2)x+x^2)(1-2tx+x^2)(1-8t^3(4t^2-5)x-10tx+x^2)}$
6	$\frac{(1+x)(x^4+80t^4x^3-24x^3t^2-2x^2-480t^4x^2+640t^6x^2+88t^2x^2+80t^4x-24t^2x+1)}{(1-x)((1+x)^2-4t^2x)((1-x)^2+16t^2(1-t^2)x)((1+x)^2-4t^2(4t^2-3)^2x)}$

Table 4. The generating function for the powers of Chebyshev polynomials of the second kind

$$\mathcal{A}_1 = a + x(b - ap),$$

$$\mathcal{A}_2 = (a^2 + xb^2)(xq - 1)a^2 + a^2p^2x(xq + 1) - 2x^2pqab,$$

$$\begin{aligned}\mathcal{A}_3 = {} & (a^3 + b^3x - a^3p^3x)(1 - q^3x^2) - 2xpq(a^3 + b^3x) - x^2a^3p^4q + 3ab^2x^2p^2q \\ & + 3ab^2x^3pq^3 - 3a^2bx^3p^2q^3 + 3a^2bx^2pq^2 - 3p^2x^2a^3q^2,\end{aligned}$$

$$\begin{aligned}\mathcal{A}_4 = {} & a^4 + (b^4 - a^4(p^4 + 3p^2q + q^2))x - q(5qa^4p^4 + b^4q + a^4q^3 + a^4p^6 + 7q^2a^4p^2 \\ & - 6qb^2a^2p^2 - 4b^3ap^3 - 4q^2ba^3p + 3b^4p^2)x^2 + q^3(-8qba^3p^3 - 3b^4p^2 + a^4q^3 \\ & + 5qa^4p^4 - 6b^2a^2p^4 - b^4q + a^4p^6 - 4q^2ba^3p + 8b^3ap^3 + 4q^2a^4p^2 + 4qb^3ap)x^3 \\ & + q^6(ap - b)^4x^4\end{aligned}$$

and

$$\mathcal{B}_1 = 1 - px - x^2q,$$

$$\mathcal{B}_2 = (1 + xq)(p^2x - (xq - 1)^2),$$

$$\mathcal{B}_3 = (1 + pqx - q^3x^2)(1 - 3pqx - p^3x - q^3x^2),$$

$$\mathcal{B}_4 = (1 - q^2x)((1 + q^2x)^2 + p^2qx)((1 - q^2x)^2 - p^2x(p^2 + 4q)).$$

Acknowledgments. The final version of this paper was written while the author was visiting University of Haifa, Israel in January 2003. He thanks the HIACS Research Center and the Caesarea Edmond Benjamin de Rothschild Foundation Institute for Interdisciplinary Applications of Computer Science for financial support, and professor Alek Vainshtein for his generosity.

REFERENCES

[HW] G.H. Hardy and E.M. Wright, An introduction to the Theory of Numbers, 4th ed. London, Oxford University Press, 1962.

[Ha] P. Haukkanen, A note on Horadam's sequence, *The Fibonacci Quarterly* **40:4** (2002) 358–361.

[Ho] A.F. Horadam, Generalization of a result of Morgado, *Portugaliae Math.* **44** (1987) 131–136.

[HM] A.F. Horadam and J.M. Mahon, Pell and Pell-Locas Polynomials, *The Fibonacci Quarterly* **23:1** (1985) 7–20.

[R] J. Riordan, Gereating function for powers of Fibonacci numbers, *Duke Math.J.* **29** (1962) 5–12.

[WR] E.T. Whittaker and G. Robinson, The Binomial Theorem, 10 in The Calculus of Observations: A Treatise on Numerical Mathematics, 4th ed., New York, Dover, 1967, 15–19.